



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Deputy Minister

Sous-ministre

MAR 02 2010

Your file / Votre référence
UNCLASSIFIED

Our file / Notre référence

MECTS # 2009-507-00175

EKME # 1170698v4

FILE # 2550-2

MEMORANDUM FOR THE MINISTER

**POTENTIAL CAUSES OF POOR RETURNS OF FRASER
RIVER SOCKEYE SALMON; WITH FOCUS ON SEA LICE IMPACTS**

(Information Only)

SUMMARY

- In a previous briefing note, ten factors were listed which could have led to the poor returns of Fraser sockeye in 2009. This note elaborates on one of those factors: the potential impacts of sea lice. Sea lice is not considered to be one of the three most likely factors but is a high profile issue.
- The most likely time when sea lice could have impacted the survival of these sockeye was in 2007, when the juvenile sockeye first entered the ocean.
- Analyses of preliminary data Fisheries and Oceans Canada (DFO) recently obtained from fish farms in the Discovery Islands region indicates that sea lice impacts likely did not contribute significantly to the low return of Fraser sockeye in 2009.
- Recently, DFO has requested sea lice and fish disease data from the fish farms operating in all areas of the BC coast.
- DFO staff will analyze these additional data to assess if either sea lice or disease from fish farms in any location in BC could have significantly contributed to the low returns of Fraser sockeye in 2009.

.../2

Background

- A previous briefing note (2009-507-00143, Tab 1) identified a number of possible factors which could have contributed to the poor returns of Fraser sockeye in 2009. These factors were:
 - Most likely – toxic algal blooms in the Strait of Georgia, low food abundance in Queen Charlotte Sound and disease;
 - Possible – predation by Humboldt squid, interception by US fisheries, sea lice from farms in Discovery Passage; and,
 - Unlikely – pollution in Fraser River, Canadian fisheries, predation on juvenile salmon in the Strait of Georgia, low food abundance in the Strait of Georgia.
- Work is continuing in Pacific Region to assess each of these factors and further information will be provided as it becomes available. Science teams have been formed to focus on each of three high profile and/or likely causes; a disease-related event, interactions with aquaculture (sea lice), and low food abundance in Queen Charlotte Sound (as well as in the Strait of Georgia).
- This note describes the potential impacts of sea lice. A previous note (MECTS 2009-507-00157, Tab 2) considered one emerging disease factor, and a forthcoming note will describe the progress on food abundance.
- Various groups including environmental organizations have speculated that sea lice from commercial fish farms are a major contributing factor to the low return of Fraser sockeye in 2009.
- In a letter to The Globe and Mail, in August 2009, Paul Sprout, Regional Director General (RDG) Pacific Region, stated: "Sea lice from fish farms are not the explanation for this year's extremely poor marine survival of Fraser River sockeye."
- More recently, one of the main conclusions from a two-day "think tank" meeting of salmon experts, organized by Simon Fraser University and the Pacific Fisheries Resource Conservation Council, was that a better understanding was still required of the potential for transmission of sea lice and disease from farmed salmon to wild salmon.
- Sea lice naturally infect all species of Pacific salmon. Historically, and long before any fish farming began in BC, most adult salmon that returned to spawn throughout BC have been infected with sea lice. Infection of the adult fish by sea lice has rarely resulted in any mortality or fish health concerns.
- The most likely time when sea lice from fish farms could have impacted the survival of the Fraser sockeye that returned in 2009 was in 2007, when the juvenile sockeye first entered the ocean.

- To examine this possibility DFO focused on fish farms in the Discovery Islands region of BC, because this area was identified by various environmental groups as the area of most concern for juvenile Fraser sockeye.
- Under BC Provincial regulations all fish farms in BC have been required since 2003 to regularly monitor and report sea lice infection levels of farmed salmon.
- Independent audits (fish sampling) of sea lice infections on all fish farms are routinely conducted by the BC Government to validate the sea lice infection data that are provided by the fish farm companies.

Analysis / DFO Comment

- With assistance from the Province of BC, DFO obtained sea lice data from the companies that operate fish farms in the Discovery Islands region.
- The data were compared for sea lice infection levels that occurred on the fish farms in the Discovery Islands from 2004-2008, for the period from April – June when most juvenile sockeye from the Fraser River were likely migrating past the fish farms in this region.
- The sea lice levels on age-1 farmed fish in 2007 were the lowest observed from 2004 - 2008.
- The sea lice levels on age-2 farmed fish in 2007 were similar to the levels that occurred in 2006, and substantially lower than the levels that occurred in 2004, 2005 and 2008.
- Overall, these results indicate that nothing unusual occurred regarding sea lice infections on the fish farms operating in the Discovery Islands region during April – June in 2007.
- While we cannot rule out the possibility that sea lice contributed to some mortality of juvenile sockeye in 2007, analysis of these preliminary data indicates that any mortality should have been similar to previous years. Thus, sea lice on fish farms operating in the Discovery Islands region likely cannot account for the unusually poor return of adult sockeye to the Fraser River in 2009.
- The sea lice data DFO used for this preliminary analyses represent the averaged results for all the fish farms that were operating in the region. However, the sea lice levels on some individual fish farms may have been higher than these averaged results indicate.

- The current DFO results are restricted to fish farms in the Discovery Islands region, and do not provide any information about sea lice levels on fish farms located in other regions of BC that might have impacted Fraser sockeye.

Recommendations / Next Steps

- DFO has requested more extensive sea lice and fish disease data from industry for all fish farms in BC.
- DFO staff will analyze these additional data to assess if either sea lice or disease from fish farms in any location in BC could have significantly contributed to the low returns of Fraser sockeye in 2009.


Claire Dansereau

Attachments

J. Kristmanson/ J. Parson/ J. Landry/ D. Gillis/ S. Paradis/ S. Mithani/ D. Bevan



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Deputy Minister

Sous-ministre

Your file Votre référence

Our file Notre référence

DEC 03 2009

MECTS#: 2009-507-00143

EKMB#: 1084753

FILE#: ABJ-8350-S2-4

MEMORANDUM FOR THE MINISTER

FACTORS AFFECTING THE 2009 FRASER SOCKEYE RETURN

(Information Only)

SUMMARY

- Sockeye salmon returns to the Fraser River in 2009 are significantly below the pre-season forecast. While the explanation for the poor 2009 Fraser return is not known, a number of factors could be important.
- Viral disease, toxic algal blooms and/or low food availability in Queen Charlotte Sound could have led to sockeye mortality at the level observed.
- Sea lice from fish farms, Humboldt squid predation and U.S. fisheries could have contributed to the sockeye mortality but are likely insufficient in themselves to explain the poor return.
- Staff continue to assemble data and analyze the key hypotheses which could inform a post-season review.

Background

- Sockeye salmon returns to the Fraser River in 2009 are significantly below the pre-season forecast. The actual return is now estimated to be on the order of 1.4 million fish, whereas more than 6 million fish¹ were expected. One exception is Harrison sockeye, which returned to the Fraser system significantly above expectation.
- Unlike other recent years when returns to the Fraser were poor, sockeye returns to Barkley Sound and the Columbia system were above expectations. However, returns to the Skeena were also poor.

.../2

¹ Fisheries and Oceans Canada provided a range of forecasts for the abundance of Fraser River sockeye in 2009, from 3.6 million to 37.6 million fish. The most commonly reported forecast is the 50% probability estimate at 10.6 million; i.e. there was a 50% probability that the return would have been greater than 10.6 million sockeye. The 75% probability estimate referenced above was 6.0 million.

- While the explanation for the poor 2009 Fraser return is not known, staff have now considered factors which could have impacted sockeye at different stages of their life cycle as they migrated from their lake-rearing habitats to the Strait of Georgia (spring/early summer 2007), on to the Gulf of Alaska and the Bering Sea and back again to spawn.

Analysis / DFO Comment

- The following factors are unlikely to have contributed to the poor 2009 return:
 1. **Pollution in the Fraser River.** There is no record of any Fraser Basin wide environmental incident that could have impacted the fish.
 2. **Capture by Canadian fisheries.** In 2009, the Canadian fishery was minimal and did not contribute to the poor return.
 3. **Predation on juvenile salmon in Strait of Georgia.** There are no known shifts in predator abundance that could explain increased predation in 2007.
 4. **Low food abundance in the Strait of Georgia.** Juvenile sockeye feed on krill. A krill fishery takes place in Jervis Inlet, but it removes a small amount of krill relative to the total krill biomass. Staff will review survey data for any evidence that juvenile sockeye were food deprived.
- The following factors may have contributed to sockeye mortality, but not at a magnitude sufficient to explain the poor return in 2009:
 1. **Predation by Humboldt squid.** Humboldt squid is a voracious predator that has increased dramatically in abundance in Canadian waters since 2007. Salmon have not been identified in their diet. Surveys in 2009 will be analyzed to assess any possible link to salmon.
 2. **Capture by U.S. fisheries.** Fraser sockeye are intercepted in U.S. Gulf of Alaska fisheries and Bering Sea fisheries. The level is not well documented but appears to be very low.
 3. **Mortality attributed to sea lice from fish farms in Discovery Passage.** While sea lice from farms could have contributed some mortality of juvenile sockeye, sea lice from natural sources could also be a factor. Staff are assessing the lice loads on farms at the time of the 2007 migration.
- The following factors could possibly have led to sockeye mortality at the scale observed:
 1. **Toxic algal blooms in the Strait of Georgia.** Extensive blooms of toxic marine algae were identified in the Strait of Georgia during 2007 when juvenile sockeye were present. Staff are working with Vancouver Island University and the aquaculture industry to assess any possible link.
 2. **Low food abundance in Queen Charlotte Sound.** Poor food supply may have impacted the survival of juvenile sockeye in Queen Charlotte Sound in the spring of 2007.

3. **Viral disease.** Preliminary evidence suggests that Fraser sockeye may be infected with a virus that could lead to mortality throughout the salmon life cycle. Staff are conducting further tests to confirm whether or not a virus could be present.

Next Steps

- Staff are continuing to assemble data and analyze the key hypotheses which could inform a post season review.
- Studies on the link to a potential viral disease are proceeding and more information is expected within the next month.


Claire Dansereau

   DEC 01 2009
NOV 27 2009



DEC 11 2009

MECTS #2009-507-00157

EKME # 1005010v8

Our file / Notre référence

MEMORANDUM FOR THE MINISTER

**POTENTIAL CAUSES OF POOR RETURNS OF FRASER RIVER
SOCKEYE SALMON: WITH FOCUS ON DISEASE-RELATED FACTORS**

(Information Only)

SUMMARY

- In a previous briefing note, ten factors were listed which could have led to the poor returns of Fraser sockeye in 2009. The three factors considered most likely to account for the scale of the mortality observed are: toxic algal blooms in the Strait of Georgia, low food abundance in Queen Charlotte Sound and disease. This note elaborates on one emerging disease factor.
- Pacific Region Science staff has documented an association between poor survival of adult migrating sockeye salmon and a particular pattern of gene responses which control physiological responses and could indicate viral activity.
- Sockeye salmon have also been found to have brain lesions. Lesions were first detected in 2009 in samples from 2006, a year when 13 million sockeye returned to the Fraser River.
- In subsequent examinations, Fisheries and Oceans Canada (DFO) staff found lesions in the brains of southern BC coho, chinook, and sockeye salmon, and across a range of life stages and sample years. The gene response pattern has also been observed across species.
- The gene response pattern and the presence of lesions have not been directly linked, but they are thought to be related to a viral infection. Molecular screening to date has not yielded a positive identification. Parasitic infections are also likely present.
- Analysis is ongoing. However, given its widespread distribution, the disease agent is thought to be endemic.
- Disease usually acts in combination with other factors (e.g., warm water conditions) rather than causing mortality directly.

.../2

Background

- In a previous briefing note (2009-507-00143, copy attached), disease was suggested as one of a number of possible factors which could have contributed to the poor returns of Fraser sockeye in 2009. These factors were:
 - Most likely – toxic algal blooms in the Strait of Georgia, low food abundance in Queen Charlotte Sound and disease,
 - Possible – predation by Humboldt squid, interception by US fisheries, sea lice from farms in Discovery Passage,
 - Unlikely – pollution in Fraser River; Canadian fisheries, predation on juvenile salmon in the Strait of Georgia, low food abundance in the Strait of Georgia.
- Work is continuing in Pacific Region to assess each of these factors and further information will be provided as it becomes available. Science teams have been formed to focus on each of three high profile and/or likely causes; a disease-related event, interactions with aquaculture (sea lice), and low food abundance in Queen Charlotte Sound (as well as in the Strait of Georgia).
- This note describes one emerging disease factor. Forthcoming notes will describe the progress on the other two.

Disease as a potential factor:

- In studies conducted over the past few years, Fisheries and Oceans Canada (DFO) staff noted evidence of poor survival in migrating adult sockeye salmon that had a particular pattern of gene response which is consistent with a viral infection. In 2006, Fraser sockeye showing this gene response experienced 30-60% higher mortality while approaching the coast and in the river than salmon that did not show the gene response pattern.
- While this pattern was first observed in samples collected from 2006, it has since been found in samples from other years back to 2003. However, a variable proportion of fish were affected each year.
- On further investigation in 2009, DFO scientists found that significant numbers of adult migrating sockeye salmon in these earlier years also contained lesions in the optic lobe of their brain and that the proportion of such fish declined sharply during the return migration. For instance, in 2009, over 70% of the sockeye salmon had lesions as they approached the coast, but this dropped to 50% in the lower river and to less than 30% at the spawning grounds. This suggests an association between the presence of lesions and en-route mortality.
- The same pattern has been found in sockeye smolts prior to them leaving the river, and in juveniles of three species (sockeye, coho, and chinook) during their first summer at sea, indicating an association with early ocean mortality in all three species.

Analysis



- The widespread observations of lesions and gene response pattern across different species, stocks, life history stages, and sample years suggest that any disease agent is endemic and has been present in salmon populations for some time.
- The gene response pattern is thought to be related to a virus. A virus from the retroviral family would be consistent with the pattern, however, few retroviruses have been described in salmon and no conclusive identification of a specific retrovirus has been made.
- The evidence of brain lesions is new and it will take some time to document the geographic extent and to understand a relationship (if any) between a disease agent and mortality.
- Other causes (than a virus) have been considered. Several different parasites are known to infect adult sockeye salmon as they migrate up the Fraser River. A myxosporean (parasitic) infection has been found (in 11 of 12 fish) in a small sample of brains of sockeye salmon so examined. Myxosporean infections are known to have a significant impact on swimming performance of sockeye smolts.
- Aquaculture operations are not considered a factor since lesions and the gene response pattern have been found in samples of sockeye smolts collected before they would have entered the marine environment.
- Other factors commonly act in combination with disease to cause mortality, rather than disease causing mortality directly. The virulence of pathogens can depend upon the level of stress of the fish (for example, those chronically stressed because of high river temperatures). Also, non-lethal infections can weaken immunity and increase the susceptibility of fish to other pathogens and to predation.
- The presence of a virus or parasite may not manifest itself as a disease or cause mortalities. In 2006, 13 million salmon returned to the Fraser River despite a high incidence of brain lesions.
- Confirmation of a disease would require isolation of the disease agent along with a demonstration of infectivity in controlled experiments.

Next Steps

- More samples are being analyzed to determine the extent of the pattern observed, for example, whether or not lesions or gene response patterns can be detected in sockeye salmon originating outside of southern BC. Additional research is also attempting to clarify which disease agents might be present.
- Further briefings on this issue will follow as necessary/requested.

- Subsequent briefings will focus on several other high profile potential causes: sea lice and low food abundance.


Claire Dansereau

Laura Richards / James Kristmanson /  Billis /  Sylvain Paradis / jr



Government of Canada
Fisheries and Oceans

Gouvernement du Canada
Pêches et Océans

MECTS # 2009-507-00175
EKME # 1170698v4
FILE / FICHER 2560-2

To: Mme. Claire Dansereau, Deputy
Pour: Minister

Date: FEB 16 2010

Object:

**POTENTIAL CAUSES OF POOR RETURNS OF FRASER
RIVER SOCKEYE SALMON: WITH FOCUS ON SEA LICE IMPACTS**

(for the Minister's Information)

From: Paul Sprout, Regional Director General, Pacific Region

FEB 16 2010

Via: Sylvain Paradis, A/Assistant Deputy Minister, Science

FEB 19 2010

Via: David Balfour, A/Assistant Deputy Minister, Fisheries & Aquaculture Mgmt.

2/2/10

Your Signature
Votre signature

Information

For Comments
Observations

Material for the Minister
Documents pour le Ministre

Remarks: L. Richards, Regional Director Science, Pacific Region
Sue Farlinger, Regional Director FAM, Pacific Region

David Balfour
A/RDS

DISTRIBUTION:

S. Johal, A/Team Leader, Executive Secretariat

Drafting Officer: Laura Richards 250-758-7177

Jim Kristmanson / Jay Parson / Jean Landry / Dave Gillis / Sylvain Paradis